

PHYSICIANS WEIGHT LOSS CLINIC

KHALED ELBANNA M.D.

Weight Loss - Aesthetic - H C G Diet

Prescription Medication Consent Form

Phentermine/Adipex: 15mg (capsule) 30mg (capsule) and 37.5 (capsule/tablet)

Indicated for patients greater than 17 years of age for short term use in conjunction with exercise and diet with the initial body mass index of $>30 \text{ kg/m}^2$, or $>27 \text{ kg/m}^2$. Works by decreasing appetite, some adverse reactions include but are not limited to dry mouth, diarrhea, and shortness of breath, chest pain, restlessness, constipation, dizziness, unpleasant taste, vomiting, heart palpitations, swollen ankles and legs, increased blood pressure, abuse dependency. Avoid abrupt withdrawal.

Phendimetrazine/Bontril: 35mg (tablet) and 105 mg ER (capsule)

Indicated for patients greater than 12 years of age for a short term use in conjunction with exercise and diet for patients with the initial body mass index of $>30\text{kg/m}^2$, or $>27\text{kg/m}^2$. This is an appetite suppressant that stimulates the central nervous system creating a chemical reaction reducing the desire to overeat. Some adverse reactions include but are not limited to diarrhea, restlessness, insomnia, dizziness, dry mouth, abdominal pain, constipation, urinary frequency, sweating, euphoria, tremors, headaches, increases, increased heart rate. Avoid abrupt withdrawal.

Diethylpropion/Tenuate: 25mg (tablet) and 75mg ER (tablet)

Indicated for patients greater than 16 years of age for short term use in conjunction with exercise and diet, with initial body mass index of $>30\text{kg/m}^2$, or $>27\text{kg/m}^2$. This is an appetite suppressant that stimulates the central nervous system creating a chemical reaction reducing the desire to overeat. Some adverse reactions include but are not limited to diarrhea, restlessness, insomnia, dizziness, dry mouth, abdominal pain, constipation, urinary frequency, sweating, euphoria, tremors, headaches, increases, increased heart rate. Avoid abrupt withdrawal.

Certain precautions should be taken for patients who are pregnant or breastfeeding, allergic to any medications or taking and OTC and cold or cough medications that may counteract the use of an appetite suppressant. Caution in mild hypertension, diabetes mellitus, and epilepsy, history of cardiovascular and pulmonary problems, and history of drug use. Notification should be given if you have any conditions or disorders that may pose a potential risk to your health while taking a medication prescribed. By signing below you agree to the consent and terms described in the above information.

Patient Name (print): _____

Patient Signature: _____ Date: _____